



ICTV Virus Taxonomy Profile: *Pospiviroidae*

Francesco Di Serio^{1,*}, Robert A. Owens², Shi-Fang Li³, Jaroslav Matoušek⁴, Vicente Pallás⁵, John W. Randles⁶, Teruo Sano⁷, Jacobus Th. J. Verhoeven⁸, Georgios Vidalakis⁹, Ricardo Flores⁵ and ICTV Report Consortium

Abstract

Members of the family *Pospiviroidae* have single-stranded circular RNA genomes that adopt a rod-like or a quasi-rod-like conformation. These genomes contain a central conserved region that is involved in replication in the nucleus through an asymmetric RNA–RNA rolling-circle mechanism. Members of the family *Pospiviroidae* lack the hammerhead ribozymes that are typical of viroids classified in the family *Avsunviroidae*. The family *Pospiviroidae* includes the genera *Apscaviroid*, *Cocadviroid*, *Coleviroid*, *Hostuviroid* and *Pospiviroid*, with >25 species. This is a summary of the ICTV Report on the family *Pospiviroidae*, which is available at ictv.global/report/pospiviroidae.

Table 1. Characteristics of members of the family *Pospiviroidae*

Example:	potato spindle tuber viroid (V01465), species <i>Potato spindle tuber viroid</i> , genus <i>Pospiviroid</i>
Genome	Single-stranded circular RNA of 246–375 nt that adopts a rod-like or quasi-rod-like conformation of minimum free energy and contains typical conserved motifs
Replication	Mediated by nuclear DNA-dependent RNA polymerase II, with oligomeric RNAs of (+) polarity cleaved by a type III RNase and circularized by DNA ligase I
Translation	Absent
Host range	Plants (dicotyledons and some monocotyledons)
Taxonomy	Several genera including >25 species

GENOME

Members of the family *Pospiviroidae* have circular single-stranded RNA genomes of a few hundred nucleotides. They may assume rod-like or quasi-rod-like conformations containing a central conserved region (CCR) and a terminal conserved hairpin (TCH) or a terminal conserved region (TCR) (Table 1, Fig. 1) [1–3]. The G+C content is >50%. The genome of viroids does not encode any proteins.

REPLICATION

Replication is nuclear and mediated by DNA-dependent RNA polymerase II, which is redirected to use RNA templates through an asymmetric RNA–RNA rolling-circle mechanism. Circular RNA molecules of (+) polarity (by convention the most abundant strand *in vivo*) are repeatedly transcribed into oligomeric complementary (–) RNAs. Such intermediates serve as templates for generating oligomeric (+) RNAs that are cleaved by a

Received 29 October 2020; Accepted 01 December 2020; Published 17 December 2020

Author affiliations: ¹Istituto per la Protezione Sostenibile delle Piante, Consiglio Nazionale delle Ricerche, Bari, 70126, Italy; ²Molecular Plant Pathology Laboratory, US Department of Agriculture, Agricultural Research Service, Beltsville, MD 20705, USA; ³State Key Laboratory of Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, 100193, PR China; ⁴Institute of Plant Molecular Biology, Biology Centre of the Czech Academy of Sciences, 37005 České Budějovice, Czech Republic; ⁵Instituto de Biología Molecular y Celular de Plantas, Universidad Politécnica de Valencia–Consejo Superior de Investigaciones Científicas, Valencia, 46010, Spain; ⁶School of Agriculture, Food and Wine, The University of Adelaide, Waite Campus, Glen Osmond, SA 5064, Australia; ⁷Faculty of Agriculture and Life Science, Hirosaki University, Hirosaki 036-8561, Japan; ⁸Plant Protection Organization of the Netherlands, Wageningen, 6700 HC, Netherlands; ⁹Department of Microbiology and Plant Pathology, University of California, Riverside, CA 92521, USA.

*Correspondence: Francesco Di Serio, francesco.diserio@ipsp.cnr.it

Keywords: *Pospiviroidae*; ICTV Report; taxonomy.

Abbreviations: CCR, central conserved region; TCH, terminal conserved hairpin; TCR, terminal conserved region.

001543 © 2021



This is an open-access article distributed under the terms of the Creative Commons Attribution License.




Fig. 1. Rod-like structure models for viroids. The positions of the central conserved region (CCR), the terminal conserved region (TCR) and the terminal conserved hairpin (TCH) are indicated by shading. The sequence-specific TCH and TCR elements have never been found together in the same viroid.

host enzyme of the RNase III class. The termini of the resulting linear monomers are ligated by the host DNA ligase 1 to generate the mature circular viroid RNA [4]. In contrast to members of the family *Avsunviroidae*, the (–) oligomeric RNAs of members of the family *Pospiviroidae* are not cleaved and do not generate the corresponding circular forms.

TAXONOMY

Current taxonomy: ictv.global/taxonomy. Demarcation of genera is based upon the type of CCR and the presence of a TCH or TCR (Fig. 1), as well as phylogenetic clustering in trees based upon whole-genome sequences (Fig. 2). Species demarcation criteria include there being <90% sequence identity and distinct biological properties with respect to other members of the genus [5]. Members of the genus *Pospiviroid*, such as potato spindle tuber viroid, share the same CCR and have a TCR. Most infect herbaceous hosts, mainly solanaceous species. Hostuviroids, such as hop stunt viroid, share the same CCR and have a TCH, except for members of the species *Dahlia latent viroid*, which have a TCR instead of the TCH. Hop stunt viroid has a wide natural host range, while dahlia latent viroid is restricted to *Dahlia* spp. Cocadviroids, such as coconut cadang-cadang viroid, share the same CCR and have a TCH. Some members infect monocotyledons, while others can only infect dicotyledons. Apscaviroids, such as apple scar skin viroid, share the same CCR and have a TCR. Apscaviroids mainly infect woody plants. Coleviroids, such as *Coleus blumei* viroid 1, share the same CCR and have a TCR or a TCH. The natural host range of coleviroids is restricted to species in the genus *Coleus*.

RESOURCES

Full ICTV Report on the family *Pospiviroidae*: ictv.global/report/pospiviroidae.




Fig. 2. Phylogenetic tree of viroid sequences. Maximum-likelihood analysis was conducted with MEGAX [6]. Nodes are labelled with bootstrap support (1000 replicates) where this was >70%.

Funding information

Production of this summary, the online chapter, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

Acknowledgements

We thank Beatriz Navarro and Michela Chiumenti for help with the phylogenetic analyses and preparation of the corresponding figures and files. Members of the ICTV Report Consortium are Stuart G. Siddell, Andrew J. Davison, Elliot J. Lefkowitz, Sead Sabanadzovic, Peter Simmonds, Donald B. Smith, Richard J. Orton and F. Murilo Zerbini.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Steger G, Perreault JP. Structure and associated biological functions of viroids. *Adv Virus Res* 2016;94:141–172.
2. Giguère T, Raj Adkar-Purushothama C, Perreault J-P. Comprehensive secondary structure elucidation of four genera of the family *Pospiviroidae*. *PLoS One* 2014;9:e98655.
3. López-Carrasco A, Flores R. Dissecting the secondary structure of the circular RNA of a nuclear viroid *in vivo*: A “naked” rod-like conformation similar but not identical to that observed *in vitro*. *RNA Biol* 2017;14:1046–1054.
4. Flores R, Gago-Zachert S, Serra P, Sanjuán R, Elena SF. Viroids: survivors from the RNA world? *Annu Rev Microbiol* 2014;68:395–414.
5. Di Serio F, Flores R, Verhoeven JTJ, Li S-F, Pallás V et al. Current status of viroid taxonomy. *Arch Virol* 2014;159:3467–3478.
6. Kumar S, Stecher G, Li M, Niyaz C, Tamara K. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 2018;35:1547–1549.